

Leptonic decay-constant ratio f_K/f_π from lattice QCD using 2+1 clover-improved fermion flavors with 2-HEX smearing

Stephan Dürr^{a,b,1}, Zoltán Fodor^{a,b,c}, Christian Hoelbling^a, Stefan Krieg^{a,b},
Laurent Lellouch^d, Thomas Lippert^{a,b}, Thomas Rae^a, Andreas Schäfer^e, Enno E. Scholz^{e,2},
Kálmán K. Szabó^{a,b}, Lukas Varnhorst^a

^a*Bergische Universität Wuppertal, Gaußstraße 20, D-42119 Wuppertal, Germany*

^b*Jülich Supercomputing Centre, Forschungszentrum Jülich, D-52425 Jülich, Germany*

^c*Institute for Theoretical Physics, Eötvös University, H-1117 Budapest, Hungary*

^d*CNRS, Aix Marseille U., U. de Toulon, CPT, UMR 7332, F-13288 Marseille, France*

^e*Regensburg University, Universitätsstraße 31, D-93053 Regensburg, Germany*

Abstract

We present a calculation of the leptonic decay-constant ratio f_K/f_π in 2+1 flavor QCD, performed via a series of lattice studies, which allows us to quantify all sources of systematic uncertainty. Our final result is $f_K/f_\pi = 1.182(28)$, and with isospin breaking corrections from ChPT the charged decay-constant ratio reads $f_{K^\pm}/f_{\pi^\pm} = 1.178(28)$. With V_{ud} from super-allowed nuclear beta decays, the CKM matrix element $V_{us} = 0.2282(54)$ is perfectly compatible with the first-row unitarity constraint of the Standard Model.

1 Introduction

Leptonic decays of pseudoscalar mesons provide a convenient way to determine Cabibbo-Kobayashi-Maskawa (CKM) matrix elements in the Standard Model (SM) and may, in the future, give access to some Beyond Standard Model (BSM) processes. Today comparison with CKM matrix elements obtained from transition form factors allows for testing the internal consistency of the SM. Obviously, in this process the highest possible precision in both theory and experiment is crucial for eventually being able to see indications for New Physics.

In this paper, we provide a computation of the decay-constant ratio f_K/f_π in the isospin symmetric limit of QCD, i.e. with two degenerate light quarks which have the same mass as the average $\frac{1}{2}(m_u^{\text{phys}} + m_d^{\text{phys}})$ in nature. By combining f_K/f_π with a factor taken from Chiral Perturbation Theory (ChPT) we determine the charged decay-constant ratio $f_{K^\pm}/f_{\pi^\pm}$. The latter object connects to the ratio of the experimentally measured widths via the relation [1]

$$\frac{\Gamma(K^\pm \rightarrow \ell \nu_\ell)}{\Gamma(\pi^\pm \rightarrow \ell \nu_\ell)} = \frac{V_{us}^2}{V_{ud}^2} \frac{f_{K^\pm}^2}{f_{\pi^\pm}^2} \frac{M_{K^\pm}}{M_{\pi^\pm}} \frac{(1 - m_\ell^2/M_{K^\pm}^2)^2}{(1 - m_\ell^2/M_{\pi^\pm}^2)^2} (1 + \delta_{\text{em}}) \quad (1)$$

¹durr (AT) itp.unibe.ch

²enno.scholz (AT) physik.uni-regensburg.de

where $\ell = e^\pm, \mu^\pm$ and ν_ℓ denotes the corresponding (electron or muon) neutrino or anti-neutrino. Here and in the following we use the standard parameterization where the complex phase in the first row of the CKM matrix is assigned exclusively to V_{ub} . Marciano advocates this form, because some of the experimental uncertainties in the determination of $\Gamma(K \rightarrow \ell \nu_\ell), \Gamma(\pi \rightarrow \ell \nu_\ell)$, some of the lattice uncertainties in the computation of f_K, f_π , and some of the radiative corrections in $\delta_{\text{em}} = \delta_K - \delta_\pi$ cancel.

Our lattice computation uses Wilson fermions [2, 3] with a tree-level clover term [4] and two levels of HEX smearing [5] along with a Symanzik improved gauge action [6]. In total we use 47 ensembles with five different lattice spacings which cover a wide range of pion masses (approximately between 130 MeV and 680 MeV) and kaon masses (such that in most cases the strange-quark mass is close to its physical value) in large boxes (such that finite-volume effects are subdominant). As we use Wilson-type fermions, a welcome feature of the Marciano setup is that the factors Z_A that would be needed to convert the bare decay constants into the physical f_π and f_K , respectively, cancel in the ratio given in Eq. (1).

These ensembles have previously been used to determine the light quark masses [7, 8], indirect CP violation [9], some low-energy constants of QCD that appear in ChPT [10], as well as the light and strange nucleon sigma terms [11]. In the present work f_K/f_π is measured on each ensemble, and the goal is to use all these data for a controlled interpolation to the physical mass point, and extrapolation to zero lattice spacing ($a \rightarrow 0$) and infinite volume ($L \rightarrow \infty$). The philosophy of the analysis is that the parameterization of the data is achieved in a modular fashion, such that we have a factor for the dependence on the quark mass, one for the dependence on the lattice spacing a , and another one for the box size L . For each factor several reasonable ansätze are considered, and the same statement holds true with respect to the cuts on the data that will be invoked (see below for details). Overall we end up with $\mathcal{O}(1000)$ reasonable analyses (i.e. combined interpolations to the physical mass point and extrapolations $a \rightarrow 0$ and $L \rightarrow \infty$). Since each one of these is performed in a fully bootstrapped fashion, we can quote a reliable estimate of both the statistical and the systematic uncertainties.

The remainder of this paper is organized as follows. In Sec. 2 we give an overview of the ensembles used in this study. We continue in Sec. 3 with a discussion of the functional ansätze which are used to establish a combined interpolation to the physical mass point and extrapolation to zero lattice spacing and to infinite volume. How these $\mathcal{O}(1000)$ analyses are distilled into a single number for f_K/f_π in the isospin limit of QCD is described in Sec. 4. This number is multiplied by a correction factor from ChPT to yield $f_{K^\pm}/f_{\pi^\pm}$, and upon combining the latter object with the PDG value $\Gamma(K \rightarrow \ell \nu_\ell)/\Gamma(\pi \rightarrow \ell \nu_\ell)$ and δ_{em} one finds V_{us}/V_{ud} as described in Sec. 5. We conclude with a check of the CKM first-row unitarity property, based on the Hardy-Towner value of V_{ud} , and compare our result to the literature.

2 Overview of ensembles used

For our analysis, we use the 2-HEX ensembles generated by the Budapest-Marseille-Wuppertal Collaboration (BMW-Collab.) with $N_f = 2 + 1$ flavors of tree-level clover-improved Wilson fermions with two HEX smearings and the tree-level Symanzik-improved gauge action [7, 8].

The two light (up- and down-) quark flavors are mass-degenerate, with their common mass m_{ud} chosen to result in pion masses between approx. 130 and 680 MeV. The single strange-quark mass m_s is taken close to the respective physical value in 45 ensembles and significantly heavier

β	a^{-1}/GeV	$\alpha_{N_f=3}$	$\alpha_{N_f=4}$
3.31	1.670(07)	0.327	0.333
3.50	2.134(15)	0.286	0.295
3.61	2.576(28)	0.262	0.271
3.70	3.031(32)	0.244	0.254
3.80	3.657(37)	0.227	0.237

Table 1: Lattice scale a^{-1} and strong-coupling parameter α for each gauge coupling β . The strong-coupling parameter α is given in the $\overline{\text{MS}}$ scheme at scale a^{-1} using both $N_f = 3$ and $N_f = 4$ matching; in our analysis we always use the average value of these two methods.

than m_s^{phys} in two ensembles. The ensembles are generated at five different gauge couplings ($\beta = 3.31, 3.50, 3.61, 3.70$, and 3.80), and this results in lattice scales a^{-1} between about 1.7 GeV and 3.7 GeV . These mass-independent lattice scales (i.e. determined for each set of ensembles at a fixed gauge coupling from the mass of the Ω -baryon at the physical mass point as described in Refs. [7, 8]) are collected in Tab. 1. In total, we have 47 different ensembles at our disposal, with particulars given in Tab. 3 of the Appendix. For more information on the ensembles and how the meson masses and decay constants are extracted from the usual two-point correlators see Refs. [7, 8, 10].

Our combination of actions is expected to result in cut-off effects which scale asymptotically like αa , where α denotes the strong coupling constant $g^2/(4\pi)$ at the scale a^{-1} (whereupon α is a logarithmic function of a). In practice, cut-off effects with similar actions are often found to scale in proportion to a^2 over the accessible range of couplings [12, 13]. This feature complicates the analysis as discussed in Sec. 3 below.

In Fig. 1 we show the combination $2M_K^2 - M_\pi^2$ of squared kaon and pion masses versus the squared pion mass M_π^2 for all of our ensembles. In this plot the y -axis serves as a somewhat non-linear representative of the simulated strange-quark mass m_s , and the x -axis serves as a slightly non-linear substitute of the joint up- and down-quark mass m_{ud} . The non-linearity in the relation between the squared meson masses and the quark mass comes from higher orders in ChPT [14]. Note that these non-linearities do not affect the definition of the physical mass point; as long as $2M_K^2 - M_\pi^2$ is a monotonic function of m_s and M_π^2 is a monotonic function of m_{ud} the requirement that $(2M_K^2 - M_\pi^2)/M_\Omega^2$ and M_π^2/M_Ω^2 would simultaneously assume their physical values leads to a unique specification of both m_s^{phys} and $m_{\text{ud}}^{\text{phys}}$.

The two panels of Fig. 2 show the kaon mass M_K and the product $M_\pi L$, respectively, as a function of the pion mass M_π . Also shown in these plots are the bounds for the fit ranges to be used in our analysis, which will be discussed in Sec. 4 below.

Regarding the definition of the physical mass point, let us add that one should not naively use the experimental values of $M_{\pi^\pm}$, $M_{K^\pm}$ and M_Ω , since these values are affected by electromagnetic self-energies and strong isospin breakings, due to $m_u^{\text{phys}} \neq m_d^{\text{phys}}$. In the FLAG report [15] it is discussed how such effects can be accounted for; here we follow the recommendation to use

$$M_\pi^{\text{phys}} = 134.8(0.3) \text{ MeV}, \quad M_K^{\text{phys}} = 494.2(0.4) \text{ MeV} \quad (2)$$

along with the PDG value $M_\Omega^{\text{PDG}} = 1.67245(29) \text{ GeV}$ from Ref. [16].




Figure 1: Scatter plot of $(2M_K^2 - M_\pi^2)$ vs. M_π^2 for all ensembles. Here the lattice scales from Tab. 1 have been used to convert the masses from lattice units into GeV. The physical mass point, as defined in Eq. (2), is indicated by a circle.

3 Overview of functional forms used

In this section we discuss the functional forms used for the fits of the decay-constant ratio in the global analysis to be presented in the following section. The functional forms have to account for the two different quark-masses that come from the joint up- and down-quark mass and the separate strange-quark mass, the different lattice volumes and the different gauge couplings (i.e. lattice scales) used in the simulations. By considering at least two different functional forms for each parameter-dependence (along with appropriate cuts), we will eventually be able to estimate the systematic uncertainties inherent in our determination of the decay-constant ratio at physical quark masses in the combined infinite volume and continuum limit.

Instead of using (renormalized) quark masses measured on the various ensembles or the bare quark mass input parameters ($am_{\text{ud}}^{\text{bare}}, am_s^{\text{bare}}$), we will always use the measured meson masses of the pion and the kaon, M_π and M_K respectively, to parameterize the quark mass dependence. In leading order ChPT, the squared pion mass is proportional to the light-quark mass, $M_\pi^2|_{\text{LO}} = 2B_0m_{\text{ud}}$, and the squared kaon mass is proportional to the sum of the two quark masses, $M_K^2|_{\text{LO}} = B_0(m_{\text{ud}} + m_s)$ [14]. Therefore, the combination $(2M_K^2 - M_\pi^2)|_{\text{LO}} = 2B_0m_s$ is in LO proportional to the strange-quark mass and can serve, beyond LO, as a non-linear substitute to parameterize the strange-quark mass dependence. As will become evident below, our analysis does not require the absence of higher-order contributions in these relations.




Figure 2: Scatter plots of M_K (left panel) and $(M_\pi L)$ (right panel) vs. M_π . Also shown are the bounds for the fit-ranges used in our analysis (see Sec. 4). Here the lattice scales from Tab. 1 have been used to convert the masses from lattice units into GeV.

In this work the dimensionless ratio f_K/f_π is considered. Nevertheless, we must decide on the scales a^{-1} of our ensembles, since we need to interpolate our data to the physical mass point and extrapolate to zero lattice spacing, $a \rightarrow 0$, and to infinite box-size, $L \rightarrow \infty$. As mentioned in the previous section, the quantity used for this purpose is the mass of the omega baryon, M_Ω . Still, there is a two-fold ambiguity regarding the scale setting procedure, and we refer to one of them as “per ensemble scale-setting” while the other one is referred to as “mass-independent scale-setting”. In both cases the lattice scale (in physical units) is obtained from

$$\frac{1}{a} = \frac{M_\Omega^{\text{PDG}}}{(aM_\Omega)} \quad (3)$$

with the PDG value of M_Ω [16]. In the former case the denominator (aM_Ω) is evaluated on each ensemble individually; this yields 47 mass-dependent scales. In the latter case the denominator is extrapolated, for each β -value, to the physical mass point (as defined in the previous section) before the relation is evaluated; this yields the five mass-independent scales listed in Tab. 1. For quantities defined at the physical mass point the “per ensemble scale-setting” and the “mass-independent scale-setting” must yield consistent results (the difference is mainly how cut-off and genuine quark mass effects are split between the respective functional ansätze). On the other hand for quantities whose definition involves derivatives with respect to quark masses the situation may be more tricky (see e.g. the discussion in Ref. [11]).

Now we turn our attention to the functional forms used to parameterize the measured decay-constant ratio. By definition the ratio f_K/f_π has to be unity along the flavor symmetric line $m_{\text{ud}} = m_s$ in the (m_{ud}, m_s) plane, where also $M_\pi = M_K$ and $f_\pi = f_K$, and we will only use

functional forms which obey the flavor-symmetry constraint

$$\left. \frac{f_K}{f_\pi} \right|_{m_{\text{ud}}=m_s} = 1 \quad (4)$$

even at finite cut-off and/or finite volume.

We start our discussion with the functional forms for the mass-dependence of the ratio f_K/f_π , from which eventually the decay-constant ratio at the physical pion- and kaon-masses will be obtained. We will either use the functional form obtained from SU(3)-ChPT up to next-to-leading order, in short NLO-SU(3) [14], or a simple polynomial expansion in the two mass-parameters. The NLO-SU(3) expression reads

$$\frac{f_K}{f_\pi} = 1 + \frac{c_0}{2} \left(\frac{5}{4} M_\pi^2 \log \left(\frac{M_\pi^2}{\mu^2} \right) - \frac{1}{2} M_K^2 \log \left(\frac{M_K^2}{\mu^2} \right) - \frac{3}{4} M_\eta^2 \log \left(\frac{M_\eta^2}{\mu^2} \right) + c_1 [M_K^2 - M_\pi^2] \right) \quad (5)$$

where we use $M_\eta^2 = (4M_K^2 - M_\pi^2)/3$, and the fit-parameters c_0, c_1 relate to the QCD low-energy constants through $c_0 = 1/(4\pi F)^2$, $c_1 = 128\pi^2 L_5(\mu)$. Throughout, μ is the renormalization scale of the chiral effective theory. In the event that c_1 or L_5 are quoted, FLAG recommends to do so for $\mu = 770 \text{ MeV}$, since this facilitates their use in phenomenology. Still we stress that the complete Eq. (5) is independent of μ . For the polynomial forms, we choose to organize the expressions in the two mass parameters M_π^2 and $M_K^2 - M_\pi^2$. This has the advantage that the flavor-symmetry constraint, Eq. (4), can be enforced through the second parameter alone, while the first parameter can account for the dominant light-quark mass-dependence. In our analysis we find that we obtain reasonable fits using these three polynomial expressions

$$\frac{f_K}{f_\pi} = 1 + [M_K^2 - M_\pi^2] \left(c_0^{3\text{-par}} + c_1^{3\text{-par}} [M_K^2 - M_\pi^2] + c_2^{3\text{-par}} M_\pi^2 \right) \quad (6)$$

$$\frac{f_K}{f_\pi} = 1 + [M_K^2 - M_\pi^2] \left(c_0^{4\text{-par}} + c_1^{4\text{-par}} [M_K^2 - M_\pi^2] + c_2^{4\text{-par}} M_\pi^2 + c_3^{4\text{-par}} M_\pi^4 \right) \quad (7)$$

$$\begin{aligned} \frac{f_K}{f_\pi} = 1 + [M_K^2 - M_\pi^2] & \left(c_0^{6\text{-par}} + c_1^{6\text{-par}} [M_K^2 - M_\pi^2] + c_2^{6\text{-par}} M_\pi^2 + c_3^{6\text{-par}} M_\pi^4 \right. \\ & \left. + c_4^{6\text{-par}} M_\pi^2 [M_K^2 - M_\pi^2] + c_5^{6\text{-par}} [M_K^2 - M_\pi^2]^2 \right) \end{aligned} \quad (8)$$

with $c_i^{3,4,6\text{-par}}$ being fit-parameters. According to the number of parameters involved these ansätze will be referred to as the 3-, 4-, and 6-parameter polynomial fits, respectively.

Next, we discuss how to parameterize the dependence on the gauge coupling (equivalently on the lattice spacing) in the simulations. Eventually, this dependence defines the continuum extrapolation $a \rightarrow 0$ of the decay-constant ratio. Given the slight difference between the asymptotically guaranteed and the observed scaling pattern that was mentioned in the previous section, it would seem natural to allow for *both* types of cut-off effects, i.e. to allow for cut-off effects proportional to $c^{\text{disc}} \alpha a + d^{\text{disc}} a^2$. As noted in previous works, with such a combined ansatz both coefficients $c^{\text{disc}}, d^{\text{disc}}$ have a tendency to be zero within errors [7, 8, 10, 11]. To account conservatively for the presence of cut-off effects in our data we chose to invoke one or the other ansatz in consecutive form. Hence we use an ansatz quadratic in the lattice spacing

$$\frac{f_K}{f_\pi} = 1 + \left(\frac{f_K}{f_\pi}(M_\pi, M_K) - 1 \right) \left(1 + c^{\text{disc}} a^2 \right) \quad (9)$$

or linear in the product αa of the strong coupling parameter and the lattice spacing

$$\frac{f_K}{f_\pi} = 1 + \left(\frac{f_K}{f_\pi}(M_\pi, M_K) - 1 \right) \left(1 + c^{\text{disc}} \alpha a \right) \quad (10)$$

to describe the discretization effects. Here the term $\frac{f_K}{f_\pi}(M_\pi, M_K)$ represents the functional form used to describe the mass-dependence, i.e. either one of Eqs. (5)–(8). Note that to obey the flavor-symmetry constraint, Eq. (4), we assume that the discretization effects only affect the deviation of the ratio from unity. The values of the strong coupling parameter α collected in Tab. 1 represent the values of α in the $\overline{\text{MS}}$ scheme evaluated at the scales given in the second column. A potential source of systematic uncertainty is that it is not clear whether one should give preference to $N_f = 4$ matching (as all of our lattice scales sitting in between m_c^{phys} and m_b^{phys} would suggest) or $N_f = 3$ matching (as would seem natural given that our ensembles are generated in QCD with $N_f = 2 + 1$). Fortunately, it turns out that the difference is too small to matter in practice; to avoid an irrelevant near-duplication of the number of analyses we simply use the average of the $N_f = 3$ and $N_f = 4$ columns. As an aside we mention that only the ratio of α at one β -value to α at another β -value matters; the whole analysis remains unchanged if the last two columns of Tab. 1 are rescaled by a common (arbitrary) factor. We emphasize that the ambiguity between the functional forms of Eqs. (9) and (10) contributes to the systematic uncertainty of our final result; if it was known for sure that the relevant cut-off effects are either $\propto \alpha a$ or $\propto a^2$, our final error-bar would be smaller.

Finally, let us discuss the volume dependence of the measured decay-constant ratio. The low-energy effective theory of QCD relates the pion mass and decay constant in finite volume $M_\pi(L), f_\pi(L)$ to the infinite volume counterparts M_π, f_π via an expansion in $\xi = M_\pi^2/(4\pi F_\pi)^2 = M_\pi^2/(8\pi^2 f_\pi^2)$ [17–20] and similar formulas are available for the kaon [20]. Unfortunately, for realistic masses and box volumes this chiral expansion is found to converge rather slowly [19, 20]. As a result of this we decided to ignore all analytical knowledge about higher-order terms and instead use the first non-trivial order version of these formulas, but with re-fitted coefficients. This means that we use the formula

$$\frac{f_K(L)}{f_\pi(L)} = \frac{f_K}{f_\pi} \left(1 + c^{\text{FV}} \left[\frac{5}{8} \tilde{g}_1(M_\pi L) - \frac{1}{4} \tilde{g}_1(M_K L) - \frac{3}{8} \tilde{g}_1(M_\eta L) \right] \right) \quad (11)$$

with the fit-parameter c^{FV} , where FV stands for finite volume, and the definitions

$$\tilde{g}_1(z) = \frac{24}{z} K_1(z) + \frac{48}{\sqrt{2}z} K_1(\sqrt{2}z) + \frac{32}{\sqrt{3}z} K_1(\sqrt{3}z) + \frac{24}{2z} K_1(2z) + \dots \quad (12)$$

$$K_1(z) = \sqrt{\frac{\pi}{2z}} e^{-z} \left\{ 1 + \frac{3}{8z} - \frac{3 \cdot 5}{2(8z)^2} + \frac{3 \cdot 5 \cdot 21}{6(8z)^3} - \frac{3 \cdot 5 \cdot 21 \cdot 45}{24(8z)^4} + \dots \right\} \quad (13)$$

where f_K/f_π represents the chosen combination of mass- and scale-dependence, as discussed previously. More terms in the expansion of $\tilde{g}_1$ are available in Ref. [19]. To have an alternative ansatz which still has the correct asymptotic behavior for large $M_\pi L$, we also use a version where both Eq. (12) and Eq. (13) are restricted to the first term. In either case the flavor-symmetry constraint, Eq. (4), is obeyed.

4 Main results obtained in the global analysis

In this section we discuss how we obtain the central value of f_K/f_π at the physical mass point, in the continuum and in infinite volume, as well as its statistical and systematic uncertainties, based on the ensembles presented in Sec. 2 and the functional forms discussed in Sec. 3.

The statistical uncertainty of a given quantity will always be determined from a bootstrap-sample of size 2000 which is generated from the bootstrap-samples of the respective underlying quantities. These bootstrap-samples will be used through every step in the fitting procedure and any further processing of the extracted parameters, in order to ensure the correct propagation of the statistical uncertainties and their correlation in every stage of our global analysis.

The systematic uncertainty, naturally, has many sources of origin. In this work we take into account the functional forms used for the mass, continuum-limit and infinite-volume extrapolations, the ranges considered in these extrapolations, and the method used for setting the lattice scale. Generally, we pursue the following strategy, which was also used in other studies, see e.g. [7, 8, 10, 11]. For each source of systematic uncertainty we consider at least two different approaches, e.g. two different methods of scale setting, and perform each fit once for each method. In that way, we arrive at many different (but valid) results for f_K/f_π at the physical mass point, in infinite volume and in the continuum. Each such result for f_K/f_π can be assigned a quality measure as provided by a suitable goodness-of-fit criterion obtained from the actual fit used (e.g. $\chi^2/\text{d.o.f.}$ or p -value). By considering the distribution (unweighted or weighted by a quality measure) of these results, we will be able to quote our final value as the mean of this distribution and determine the systematic uncertainty from its width (variance). If we consider a subset of these results, e.g. the subset generated with one specific functional ansatz for the volume dependence, the resulting distribution will be narrower than the full distribution. In this way we can provide an error budget, i.e. we can break up the full systematic error into individual contributions.

For the mass dependence, we consider four different functional forms, Eqs. (5–8), which have two, three, four, or six fit parameters, respectively. For the continuum extrapolation we either use the “discretization via a^2 ” ansatz, Eq. (9), or the “discretization via αa ” ansatz, Eq. (10), both of which have one fit parameter. The infinite volume extrapolation uses the functional form of Eq. (11), where either the full $\tilde{g}_1$ function is used or only its leading term, invoking one fit parameter in both cases. This already leads to $4 \times 2 \times 2 = 16$ different combinations of functional forms used in the extrapolation of the ratio f_K/f_π , which (depending on the mass extrapolation considered) have either four, five, six, or eight fit parameters in total. Finally, the number of fit ansätze is doubled due to the two methods of setting the scale (“mass-independent” versus “per-ensemble”, see Sec. 3). Therefore we shall consider 32 different combinations of interpolation/extrapolation and scale setting methods in our global analysis.

Another source of spread in the final distribution is the choice of ensembles used in the global analysis. In principle, one could consider all $2^{47} = \mathcal{O}(10^{14})$ combinations that result from including or excluding any one of the 47 available ensembles. Since this is neither sensible nor feasible, we restrict ourselves to just a few cuts in the pion- and kaon-mass, the volume parameter ($M_\pi L$), and the range of lattice scales or gauge couplings β . For the pion mass M_π of the selected ensembles we use either no upper bound, or $M_\pi^{\text{max}} = 350 \text{ MeV}$, 300 MeV , or 250 MeV , whereas no lower bound is used (i.e. the ensembles with nearly physical pion mass are always included, unless there is another restriction). Also for the kaon mass M_K no lower bound is used, and the upper bounds M_K^{max} are: none, 600 MeV , 550 MeV , 500 MeV .

For the volume extrapolation, we consider either no minimal bound for the parameter $(M_\pi L)$ or $(M_\pi L)^{\min} = 3.85$ or 4.05 . For the readers convenience these choices are indicated in the plots of Fig. 2 by horizontal and vertical dashed lines. For the gauge coupling β we consider the lower bounds $\beta \geq \beta^{\min} = 3.31, 3.50$ and 3.61 and no upper bound. Additionally, we only consider those combinations of fit ranges which contain at least five ensembles, since the minimal number of fit parameters in our fits is already four. By excluding double counting of fit ranges (combination of bounds which lead to the same set of included ensembles) we arrive at 63 possible sets of fit ranges.

In conjunction with the 32 combinations of extrapolations one might expect that this number of 63 fit ranges would lead to $32 \times 63 = 2,016$ different fits. But one has to take into account that some combinations of fit ranges do not include sufficiently many ensembles to allow for fit functions with five, six, or eight parameters. In our analysis we shall only consider “true fits”, that is fits which have at least one degree of freedom. Also accounting for the rare cases in which the used fitting routine would not find a unique minimum, we arrive at a total of 1368 single fits used in the global analysis.

To give the reader an impression of how these fits work out in practice, we include a total of five plots. In Fig. 3 a scatter-plot of the p -value (as a goodness-of-fit measure obtained in each fit) versus the final decay-constant ratio is shown. The grey band indicates the overall uncertainty (statistical and systematic errors added in quadrature) of our final result as specified in Eq. (16) below. There is a number of low-order polynomial fits with surprisingly good p -values to the left of the grey band; this explains why the weighted average is smaller than the unweighted average (see below).

More details for two examples from our 1368 fits are shown in Figs. 4 and 5. They use the SU(3)-ChPT functional form and the 4-parameter polynomial form, respectively, for the mass-dependence. Both fits were performed with the a^2 -discretization ansatz, with full FV-dependence and with the mass-independent scale setting method. The SU(3)-ChPT fit shown in Fig. 4 includes only ensembles with pion masses below 300 MeV, kaon masses below 600 MeV, along with $(M_\pi L) \geq 3.85$, and $\beta \geq 3.50$. The 4-parameter polynomial fit shown in Fig. 5 includes all ensembles with $\beta \geq 3.61$. In general, we find that polynomial fits with suitable restrictions in the meson masses and volumes tend to give good fits (as judged by the p -values), but they show a wildly fluctuating behavior outside of the admitted region, especially above the selected pion mass cut. Evidently, this behavior is neither unexpected nor does it invalidate using results extracted at the physical mass point from such a fit, since the physical meson mass region is always included in the fit range. Perhaps the main difference between the fit shown in Fig. 4 and the one shown in Fig. 5 is that the former one (with good χ^2) describes the data outside of the fit interval (represented by black symbols in the right panel) quite well, whereas the latter one (with poor χ^2) misses the data outside of its fit range quite visibly.

The final step is to perform an average of our 1368 fits, each of which has a central value and a statistical uncertainty. This average may be performed with a uniform weight for all fits or it may be performed in a weighted manner, and a similar statement holds for the standard deviation of the 1368 fitted values, which we use as a measure of the systematic uncertainty of the aggregate f_K/f_π . Regarding the weights, it is natural to consider weights which derive from some goodness-of-fit number of the individual fits. One option is to use a weight proportional to the p -value of each fit. In Tab. 2 we collect the various averages that result from our 1368 valid analyses with a flat weight in the left column and with the p -value weight in the right column. In addition, the resulting aggregate values (along with their statistical and systematic




Figure 3: Scatter plot of the p -value of the fit vs. the physical-world ratio f_K/f_π from each one of the 1368 single fits used in the global analysis. The symbols distinguish fits using the ChPT or the polynomial form for the interpolation in the quark masses. Additionally, for the results using the polynomial form, the color indicates whether the 3-, 4-, or 6-parameter polynomial form has been used. For better readability no error-bars are shown for the points in this figure. The vertical *solid* and *dashed* lines and *grey band* show our final result from Eq. (16) and its statistical and total uncertainty, respectively.

uncertainties) are shown for all the data cuts considered and for all functional forms employed; this amounts to a break-up of the overall uncertainty into its various contributions. Instead of using the p -value as a weight, we tried alternatives such as the weights based on the Akaike information criterion [21] or based on $\chi^2/n_{\text{d.o.f.}}$. Basically, we find that using these weights does not change the mean values and error estimates significantly compared to the results obtained with the p -value, and this is why we refrain from including them in the table.

A graphical summary of the content of Tab. 2 is shown in Fig. 6. The black vertical line indicates our central value as given in Eq. (16) below, the dashed set of vertical lines shows the statistical uncertainty, and the grey band displays the combined uncertainty as specified in that equation. For each fit ansatz (left panel) and cut on the data (right panel) the respective entry in Tab. 2 is shown, enabling one to spot trends (e.g. in β^{min}) more easily.

To summarize Tab. 2, we note that from the unweighted and p -value weighted distributions




Figure 4: Example of a fit with SU(3)-ChPT mass-dependence, a^2 -discretization, full FV, and mass-independent a^{-1} . Included are ensembles with $M_\pi \leq 300$ MeV, $M_K \leq 600$ MeV, $(M_\pi L) \geq 3.85$, and $\beta \geq 3.50$. The *left panel* compares the value of f_K/f_π from the fit and measured on the ensemble for each data point (labeled from 0 to 46, ordered as in Tab. 3). The *right panel* shows the dependence of the fitted f_K/f_π on the squared pion mass M_π^2 at the physical strange-quark mass and in the infinite volume and continuum limit. The data-points shown in this panel have been shifted according to these values/limits using the fit results. This particular fit resulted in $\chi^2 = 7.6$ with $n_{\text{d.o.f.}} = 6$, giving a p -value of 0.27.

we obtain from all fits the final results

$$f_K/f_\pi|_{\text{flat}} = 1.191(08)_{\text{stat}}(24)_{\text{syst}} = 1.191(25)_{\text{comb}}, \quad (14)$$

$$f_K/f_\pi|_{p\text{-value}} = 1.173(11)_{\text{stat}}(29)_{\text{syst}} = 1.173(31)_{\text{comb}}, \quad (15)$$

where the combined error was obtained by adding the statistical and systematic errors in quadrature. The only question left open is to which one of the two weighting strategies preference should be given. In the end, we decided to take a straight average, resulting in

$$f_K/f_\pi = 1.182(10)_{\text{stat}}(26)_{\text{syst}} = 1.182(28)_{\text{comb}} \quad (16)$$

as our final result for the decay-constant ratio in $N_f = 2 + 1$ QCD at the physical mass point in the combined infinite volume and continuum limit.

5 Discussion

The decay-constant ratio in Eq. (16) represents our final result for QCD with two degenerate flavors taken at the average mass of the up- and down-quarks in the real world and a single flavor taken at the physical strange-quark mass. Its systematic error includes all sources of




Figure 5: Example of a fit with 4-parameter polynomial mass-dependence, a^2 -discretization, full FV, and mass-independent a^{-1} . Included are ensembles with $\beta \geq 3.61$. The *left panel* compares the value of f_K/f_π from the fit and measured on the ensemble for each data point (labeled from 0 to 46, ordered as in Tab. 3). The *right panel* shows the dependence of the fitted f_K/f_π on the squared pion mass M_π^2 at the physical strange-quark mass and in the infinite volume and continuum limit. The data-points shown in this panel have been shifted according to these values/limits using the fit results. This particular fit resulted in $\chi^2 = 72.1$ with $n_{\text{d.o.f.}} = 18$, giving a p -value of $10^{-8} \approx 0$.

theoretical uncertainty in that theory. However, the quantity we are after is $f_{K^\pm}/f_{\pi^\pm}$ in the real world, i.e. in QCD with six non-degenerate flavors, each of which to be taken at its own physical mass. This change will bring a shift of the central value, as well as an increase in the systematic uncertainty, and our goal is to discuss these effects.

Regarding the influence of those quark flavors which were ignored in Eq. (16) it is clear that the dominant unquenching effect comes from the lightest flavor ignored, i.e. the charm quark. Since the QCD functional determinant is quadratic in the masses of the quarks, naive reasoning would suggest that unquenching effects from charm quark loops are suppressed, relative to those from strange quarks, by a factor $(m_c/m_s)^2 \simeq 140$. Since already the presence of strange quark loops in the sea seems to affect f_K/f_π very little [15], we take it for granted that the quenching of the heavier flavors introduces an uncertainty which is small in comparison to the one which is declared in Eq. (16) and can thus be considered negligible in what follows.

The correction needed to undo the isospin symmetry limit in Eq. (16) is also small compared to the uncertainty in that equation. However, the issue has been analyzed in ChPT [14, 22], relating the object of interest, $f_{K^\pm}/f_{\pi^\pm}$, to the isospin symmetric quantity f_K/f_π through

$$\frac{f_{K^\pm}}{f_{\pi^\pm}} = \frac{f_K}{f_\pi} \sqrt{1 + \delta_{\text{SU}(2)}} \quad (17)$$

but the information on $\delta_{\text{SU}(2)}$ in the literature is not very conclusive. The original work suggests

fit type	flat	p -value	valid fits
all	1.191(08)(24)	1.173(11)(29)	1368
$M_\pi^{\max} = 250$ MeV	1.194(18)(29)	1.180(21)(43)	104
$M_\pi^{\max} = 300$ MeV	1.184(12)(22)	1.168(15)(28)	222
$M_\pi^{\max} = 350$ MeV	1.196(09)(23)	1.177(11)(21)	380
$M_\pi^{\max} = \infty$	1.189(07)(24)	1.171(10)(25)	662
$M_K^{\max} = 500$ MeV	1.190(27)(12)	1.197(33)(11)	12
$M_K^{\max} = 550$ MeV	1.182(11)(18)	1.174(12)(26)	396
$M_K^{\max} = 600$ MeV	1.198(09)(23)	1.168(13)(32)	680
$M_K^{\max} = \infty$	1.186(06)(28)	1.188(13)(23)	280
$(M_\pi L)^{\min} = 0$	1.191(10)(21)	1.169(15)(33)	623
$(M_\pi L)^{\min} = 3.85$	1.191(10)(24)	1.175(14)(23)	497
$(M_\pi L)^{\min} = 4.05$	1.188(10)(31)	1.179(16)(28)	248
$\beta^{\min} = 3.31$	1.194(08)(17)	1.193(12)(20)	682
$\beta^{\min} = 3.50$	1.183(09)(17)	1.168(13)(22)	450
$\beta^{\min} = 3.61$	1.195(14)(42)	1.135(23)(25)	236
scale setting mass-independent	1.188(08)(27)	1.168(12)(32)	719
scale setting per-ensemble	1.194(08)(20)	1.179(12)(24)	649
discretization via a^2 -term	1.189(07)(24)	1.169(12)(31)	700
discretization via αa -term	1.193(09)(24)	1.178(11)(24)	668
finite volume via full $\tilde{g}_1$ function	1.191(08)(24)	1.172(11)(29)	688
finite volume via leading terms	1.191(08)(24)	1.174(11)(28)	680
quark mass function via ChPT	1.189(07)(21)	1.183(10)(19)	341
quark mass function via 3-par. poly.	1.181(08)(26)	1.159(14)(30)	382
quark mass function for 4-par. poly.	1.198(09)(21)	1.183(13)(27)	367
quark mass function for 6-par. poly.	1.197(09)(22)	1.177(11)(27)	278

Table 2: Comparison of the unweighted (“flat”) and p -value weighted results from the complete set of fits (top line) with the various subsets (according to $M_\pi^{\max}$, $M_K^{\max}$, $(M_\pi L)^{\min}$, $\beta^{\min}$, the scale setting method, the type of discretization terms, the type of finite volume terms, and the kind of quark mass dependence used) which allows for a breakup of the systematic uncertainty into various sources (see text for details). The first error is statistical and the second one is systematic. The last column gives the number of single fits which match these criteria.

$\delta_{\text{SU}(2)} = -0.0043(12)$ [22] and FLAG finds, from a re-analysis of several $N_f = 2+1$ computations, a very similar value [15]. On the other hand, a dedicated study in $N_f = 2$ QCD ends up finding $\delta_{\text{SU}(2)} = -0.0078(7)$ [23]. In this situation we opt for $\delta_{\text{SU}(2)} = -0.0061(61)_{\text{syst}}$ with a generous 100% error. Upon combining Eq. (16) with this estimate of SU(2) breaking effects, we find

$$f_{K^\pm}/f_{\pi^\pm} = 1.178(10)_{\text{stat}}(26)_{\text{syst}} = 1.178(28)_{\text{comb}} \quad (18)$$

which is the quantity to be used in the final phenomenological analysis.

The next ingredient is an evaluated form of the original Marciano relation, Eq. (1), with all uncertainties adequately propagated. In the literature we find [24, 25]

$$\frac{V_{\text{us}}}{V_{\text{ud}}} \frac{f_{K^\pm}}{f_{\pi^\pm}} = 0.27599(29)(24) = 0.27599(38)_{\text{exp}} \quad (19)$$




Figure 6: Overview of the results from the global analysis with different fit types (*left*) and different fit ranges (*right*). Shown are the value and the statistical and combined statistical and systematic errors, see Tab. 2 for details. The vertical black solid and dashed lines and the grey shaded band show the central value, the statistical and combined error, respectively, of our final result for f_K/f_π in 2+1 flavor QCD as given in Eq. (16).

which, when combined with our result (18), yields

$$\frac{V_{\text{us}}}{V_{\text{ud}}} = 0.2343(20)_{\text{stat}}(52)_{\text{syst}}(03)_{\text{exp}} = 0.2343(55)_{\text{comb}} \quad (20)$$

where “stat” and “syst” refer to our statistical and systematic uncertainties, respectively, while “exp” refers to the combined uncertainty of Eq. (19), which is comparatively small.

With this ratio of CKM matrix elements in hand, and given that $|V_{\text{ub}}| = 4.12(37)(06) \cdot 10^{-3}$ [25] is small on the scale of our uncertainties, one can proceed in two ways. One option is to assume the unitarity inherent in the CKM paradigm. In this case the square of Eq. (20) is augmented with the first-row unitarity relation in the form $V_{\text{ud}}^2 \cdot (1 + V_{\text{us}}^2/V_{\text{ud}}^2) = 0.999983(3)$ to yield

$$V_{\text{ud}} = 0.9736(04)_{\text{stat}}(11)_{\text{syst}}(01)_{\text{exp}} = 0.9736(12)_{\text{comb}} , \quad (21)$$

while the inverse square of Eq. (20) is combined with the first-row unitarity relation in the form $V_{\text{us}}^2 \cdot (1 + V_{\text{ud}}^2/V_{\text{us}}^2) = 0.999983(3)$ to yield

$$V_{\text{us}} = 0.2281(18)_{\text{stat}}(48)_{\text{syst}}(03)_{\text{exp}} = 0.2281(51)_{\text{comb}} . \quad (22)$$

The other option is to multiply the very precise result from super-allowed nuclear beta decays, $V_{\text{ud}} = 0.97417(21)_{\text{nuc}}$ by Hardy and Towner [26], with our result Eq. (20) to give

$$V_{\text{us}} = 0.2282(19)_{\text{stat}}(51)_{\text{syst}}(03)_{\text{exp+nuc}} = 0.2282(54)_{\text{comb}} , \quad (23)$$

and in the same way one may form the product $V_{\text{ud}}^2 \cdot (1 + V_{\text{us}}^2/V_{\text{ud}}^2)$ with the Hardy-Towner value as the first factor and our result Eq. (20) as the second factor and add $|V_{\text{ub}}|^2$ to find

$$V_{\text{ud}}^2 + V_{\text{us}}^2 + |V_{\text{ub}}|^2 - 1 = 0.0011(09)_{\text{stat}}(23)_{\text{syst}}(05)_{\text{exp+nuc}} = 0.0011(25)_{\text{comb}} \quad (24)$$

which indicates that the unitarity relation is well obeyed, within errors.

6 Summary

In this paper we have presented a lattice computation of f_K/f_π in 2+1 flavor QCD, i.e. in the isospin limit with the two light quarks taken at the average of the physical up- and down-quark masses, while the non-degenerate last flavor is taken at the physical value of the strange-quark mass. The final value in Eq. (16) gives the combined continuum and infinite volume limit, and it accounts for all sources of systematic uncertainty in that theory.

In the next step we have used external information to account for the breaking of isospin symmetry in the real world. Fortunately, the correction is very small, and our result for $f_{K^\pm}/f_{\pi^\pm}$ as given in Eq. (18) has the same error-bar as its isospin symmetric counterpart. Still, when comparing to the literature we find that our uncertainty is fairly large. The MILC value $f_{K^\pm}/f_{\pi^\pm} = 1.1947(26)(37)$ [27], the HPQCD value $f_{K^\pm}/f_{\pi^\pm} = 1.1916(15)(15)$ [28], the FNAL/MILC value $f_{K^\pm}/f_{\pi^\pm} = 1.1956(10)(^{+26}_{-18})$ [29] and the ETMC value $f_{K^\pm}/f_{\pi^\pm} = 1.184(12)(11)$ [30] all stem from simulations with $N_f = 2 + 1 + 1$ dynamical flavors, and concern the charged decay-constant ratio. The RBC/UKQCD value $f_K/f_\pi = 1.1945(45)$ [31] comes from simulations with $N_f = 2 + 1$ flavors of domain-wall fermions, and requires the same (small) isospin breaking correction that we have applied to our result. What catches our attention is that the overall uncertainties quoted by ETMC and in our Eq. (18) are roughly an order of magnitude larger than the overall uncertainties obtained with staggered and domain-wall fermions.

In a last step we have explored the implications of our result for the charged decay-constant ratio on the first row of the CKM matrix. By invoking only experimental information we find Eq. (20). If we assume that the CKM matrix is unitary, we find the individual elements Eq. (21) and Eq. (22), respectively. While not very precise, they are at least consistent with the averages given in Refs. [15, 25]. Alternatively, we may take the Hardy-Towner value of V_{ud} as an additional input. In this case we can test whether our result is consistent with the CKM matrix being unitary, and Eq. (24) tells us that, within errors, this is the case.

Acknowledgments

Computations were carried out on the BG/Q supercomputer JUQUEEN at Forschungszentrum Jülich through a NIC grant, on Turing at IDRIS, France, under GENCI-IDRIS grant 52275, and on a local cluster at the University of Wuppertal. This work was supported by the DFG through the SFB/TRR 55 “Hadron Physics from Lattice QCD”, by the EU Framework Programme 7 grant (FP7/2007-2013)/ERC No 208740, by the OTKA grant OTKA-NF-104034, and by the projects OCEVU Labex (ANR-11-LABX-0060) and A*MIDEX (ANR-11-IDEX-0001-02). In addition, E.E.S. acknowledges support from the EU grant PIRG07-GA-2010-268367.

Appendix: Ensemble details

In this appendix we arrange some details on the ensembles used in this work. To simulate the $N_f = 2 + 1$ flavors with the clover-improved fermion action with 2-HEX smearing and the Symanzik-improved gauge action, the Hybrid-Monte-Carlo (HMC) algorithm has been used together with the rational approximation variety (RHMC) to represent the single strange quark flavor. More details on the implementation of these algorithms can be found in Ref. [8].

β	am_{ud}^{bare}	am_s^{bare}	L/a	aM_π	aM_K	aM_Ω	f_K/f_π
3.31	-0.07000	-0.0400	16	0.35262(104)	0.40757(96)	1.1364(189)	1.0478(18)
	-0.09000	-0.0400	24	0.20838(76)	0.33265(64)	1.0408(90)	1.1125(30)
	-0.09000	-0.0440	24	0.20266(121)	0.31998(95)	1.0327(144)	1.1089(58)
	-0.09300	-0.0400	24	0.17789(57)	0.32029(40)	1.0126(47)	1.1359(33)
	-0.09300	-0.0400	32	0.17697(51)	0.32001(42)	1.0212(52)	1.1281(35)
	-0.09300	-0.0440	32	0.17182(112)	0.30812(83)	0.9912(107)	1.1258(90)
	-0.09530	-0.0400	32	0.14967(74)	0.30985(65)	1.0141(81)	1.1354(82)
	-0.09530	-0.0440	32	0.14534(120)	0.29809(75)	1.0014(114)	1.1660(97)
	-0.09756	-0.0400	32	0.11928(98)	0.30052(59)	0.9898(60)	1.1727(85)
	-0.09900	-0.0400	48	0.08950(69)	0.29352(35)	0.9861(36)	1.1887(56)
	-0.09933	-0.0400	48	0.08111(107)	0.29257(87)	0.9918(110)	1.1974(198)
3.50	-0.02500	-0.0060	16	0.28911(79)	0.32587(72)	0.8983(165)	1.0433(08)
	-0.03100	-0.0060	24	0.25373(45)	0.30566(43)	0.8654(85)	1.0578(15)
	-0.03600	-0.0060	24	0.22506(71)	0.29175(62)	0.8553(105)	1.0716(26)
	-0.04100	-0.0060	24	0.19249(46)	0.27697(41)	0.8332(37)	1.1008(14)
	-0.04100	-0.0120	24	0.18851(77)	0.26121(71)	0.8058(146)	1.0961(42)
	-0.04370	-0.0060	24	0.17379(45)	0.26928(37)	0.8269(33)	1.1205(24)
	-0.04630	-0.0120	32	0.14399(59)	0.24358(59)	0.8051(100)	1.1210(46)
	-0.04800	-0.0023	32	0.13538(56)	0.26494(55)	0.8297(59)	1.1631(62)
	-0.04900	-0.0060	32	0.12089(83)	0.25090(70)	0.8104(77)	1.1679(84)
	-0.04900	-0.0120	32	0.11792(56)	0.23549(46)	0.7937(64)	1.1458(76)
	-0.05150	-0.0120	48	0.08472(50)	0.22578(62)	0.7613(78)	1.1846(107)
	-0.05294	-0.0060	64	0.06121(62)	0.23578(65)	0.7877(49)	1.2211(147)
3.61	-0.02000	-0.0042	32	0.19645(33)	0.23343(32)	0.7020(60)	1.0503(08)
	-0.02000	-0.0045	32	0.19889(29)	0.25382(28)	0.7204(51)	1.0724(10)
	-0.02800	0.0045	32	0.14855(42)	0.23381(36)	0.7051(48)	1.1249(26)
	-0.03000	-0.0042	32	0.12947(46)	0.20528(43)	0.6642(53)	1.1126(32)
	-0.03000	0.0045	32	0.13221(39)	0.22734(34)	0.6926(43)	1.1441(35)
	-0.03121	0.0045	32	0.12094(23)	0.22398(25)	0.6986(40)	1.1456(28)
	-0.03300	0.0045	48	0.10271(46)	0.21834(47)	0.6786(49)	1.1629(61)
	-0.03440	0.0045	48	0.08588(46)	0.21398(48)	0.6828(49)	1.1997(74)
	-0.03650	-0.0030	64	0.04713(43)	0.18632(52)	0.6383(54)	1.2280(103)
3.70	-0.00500	0.0500	32	0.22281(38)	0.32097(39)	0.7952(54)	1.1268(30)
	-0.01500	0.0000	32	0.16439(37)	0.20127(36)	0.6128(116)	1.0582(19)
	-0.01500	0.0500	32	0.17104(35)	0.30039(35)	0.7943(33)	1.1845(29)
	-0.02080	-0.0050	32	0.12498(38)	0.17172(34)	0.5653(97)	1.0912(28)
to be continued on next page							

β	$am_{\text{ud}}^{\text{bare}}$	$am_{\text{s}}^{\text{bare}}$	L/a	aM_{π}	aM_K	aM_{Ω}	f_K/f_{π}
continued from previous page							
	-0.02080	0.0000	32	0.12464(51)	0.18368(47)	0.5841(113)	1.1007(36)
	-0.02080	0.0010	32	0.12514(46)	0.18675(45)	0.5678(68)	1.1177(38)
	-0.02540	-0.0050	48	0.08035(30)	0.15519(28)	0.5392(55)	1.1348(54)
	-0.02540	0.0000	48	0.08166(29)	0.16987(30)	0.5585(43)	1.1606(54)
	-0.02700	0.0000	64	0.06038(32)	0.16376(33)	0.5569(57)	1.1649(68)
3.80	-0.00900	0.0000	32	0.15246(35)	0.17394(33)	0.5116(94)	1.0453(14)
	-0.01400	0.0000	32	0.12053(57)	0.15893(51)	0.5096(122)	1.0908(32)
	-0.01400	0.0030	32	0.12304(54)	0.16818(52)	0.4797(64)	1.0939(33)
	-0.01900	0.0000	48	0.08200(86)	0.14461(90)	0.4626(43)	1.1398(127)
	-0.01900	0.0030	48	0.08230(105)	0.15382(64)	0.4743(65)	1.1819(145)
	-0.02100	0.0000	64	0.05984(22)	0.13947(39)	0.4658(33)	1.1657(113)

Table 3: Simulated ensembles and measured aM_{π} , aM_K , aM_{Ω} , and f_K/f_{π} .

The above table provides information about the individual ensembles for one choice of the extraction details. The first four columns contain the input parameters for each simulation, i.e. the gauge coupling β , the values of the bare mass parameters $am_{\text{ud}}^{\text{bare}}$, $am_{\text{s}}^{\text{bare}}$ for the light and strange quarks, respectively, and the spatial extent in lattice units L/a (the temporal extent of the lattices is typically larger and not listed here). Note that the bare mass parameters can be negative, due to the additive quark mass renormalization with Wilson-type fermions, and the resulting (renormalized) quark masses are still positive. The remaining columns show the quantities measured on these ensembles which are relevant for this work, and their statistical uncertainty as determined from the bootstrap procedure. We like to stress that this table does not show the information on the correlation between measurements on the same ensembles, but of course by using the bootstrap samples for each quantity these correlations are properly taken into account in our analysis. Further details on how these quantities were extracted can be found in Ref. [10].

References

- [1] W. J. Marciano, Phys. Rev. Lett. **93**, 231803 (2004) [hep-ph/0402299].
- [2] K.G. Wilson, Phys. Rev. D **10**, 2445 (1974).
- [3] K.G. Wilson, *New Phenomena In Subnuclear Physics. Part A. Proceedings of the First Half of the 1975 International School of Subnuclear Physics, Erice, Sicily, July 11 - August 1, 1975*, ed. A. Zichichi, Plenum Press, New York, 1977, p. 69, CLNS-321.
- [4] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B **259**, 572 (1985).
- [5] S. Capitani, S. Dürr and C. Hoelbling, JHEP **0611**, 028 (2006) [hep-lat/0607006].
- [6] M. Lüscher and P. Weisz, Phys. Lett. B **158**, 250 (1985).
- [7] S. Dürr *et al.*, Phys. Lett. B **701** (2011) 265 [arXiv:1011.2403].
- [8] S. Dürr *et al.*, JHEP **1108** (2011) 148 [arXiv:1011.2711].
- [9] S. Dürr *et al.*, Phys. Lett. B **705**, 477 (2011) [arXiv:1106.3230].
- [10] S. Dürr *et al.* [Budapest-Marseille-Wuppertal Collaboration], Phys. Rev. D **90** (2014) 11, 114504 [arXiv:1310.3626].
- [11] S. Dürr *et al.*, arXiv:1510.08013 [hep-lat].
- [12] T.A. DeGrand, A. Hasenfratz and T.G. Kovacs [MILC Collaboration], hep-lat/9807002.
- [13] S. Dürr, Z. Fodor, C. Hoelbling, R. Hoffmann, S. D. Katz, S. Krieg, T. Kurth and L. Lelouch *et al.*, Phys. Rev. D **79**, 014501 (2009) [arXiv:0802.2706].
- [14] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 465.
- [15] S. Aoki *et al.*, Eur. Phys. J. C **74** (2014) 2890 [arXiv:1310.8555].
- [16] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38** (2014) 090001.
- [17] M. Lüscher, Commun. Math. Phys. **104**, 177 (1986).
- [18] J. Gasser and H. Leutwyler, Phys. Lett. B **184**, 83 (1987).
- [19] G. Colangelo and S. Dürr, Eur. Phys. J. C **33**, 543 (2004) [hep-lat/0311023].
- [20] G. Colangelo, S. Dürr and C. Haefeli, Nucl. Phys. B **721**, 136 (2005) [hep-lat/0503014].
- [21] H. Akaike. 1974. IEEE Transactions on Automatic Control 19, 716.
- [22] V. Cirigliano and H. Neufeld, Phys. Lett. B **700**, 7 (2011) [arXiv:1102.0563].
- [23] G. M. de Divitiis *et al.*, JHEP **1204**, 124 (2012) [arXiv:1110.6294].
- [24] M. Moulson, arXiv:1411.5252 [hep-ex].
- [25] J. L. Rosner, S. Stone and R. S. Van de Water, arXiv:1509.02220 [hep-ph].
- [26] J. C. Hardy and I. S. Towner, Phys. Rev. C **91**, no. 2, 025501 (2015) [arXiv:1411.5987].
- [27] A. Bazavov *et al.* [MILC Collaboration], Phys. Rev. Lett. **110**, 172003 (2013) [arXiv:1301.5855].
- [28] R. J. Dowdall, C. T. H. Davies, G. P. Lepage and C. McNeile [HPQCD Collaboration], Phys. Rev. D **88**, 074504 (2013) [arXiv:1303.1670].
- [29] A. Bazavov *et al.* [Fermilab Lattice and MILC Collaborations], Phys. Rev. D **90**, 074509 (2014) [arXiv:1407.3772].
- [30] N. Carrasco *et al.*, Phys. Rev. D **91**, 054507 (2015) [arXiv:1411.7908].
- [31] T. Blum *et al.* [RBC and UKQCD Collaborations], arXiv:1411.7017 [hep-lat].